

## Nonbook problems

1. Verify that for any nonzero constant  $b$ , the function  $f(x) = \frac{1}{b} \cosh(bx)$  satisfies the differential equation

$$\frac{d^2y}{dx^2} - b\sqrt{1 + \left(\frac{dy}{dx}\right)^2} = 0.$$

(Recall that the function "cosh" is defined by  $\cosh(x) = \frac{1}{2}(e^x + e^{-x})$ .)

2. Verify that on the interval  $-2 < x < 2$ , the two continuous functions  $y(x) = \sqrt{4 - x^2}$  and  $y(x) = -\sqrt{4 - x^2}$ , obtainable from the implicit solution  $x^2 + y^2 = 4$  of the differential equation  $x + y \, dy/dx = 0$ , are (explicit) solutions of this differential equation.

3. Show that if  $L_1$  and  $L_2$  are linear operators, then  $L_1 + L_2$  is a linear operator.

4. Show by induction on  $n$  that if an operator  $L$  is linear, then for all  $n \geq 1$ , all constants  $c_1, c_2, \dots, c_n$ , and all functions  $f_1, f_2, \dots, f_n$ ,

$$L[c_1f_1 + c_2f_2 + \dots + c_nf_n] = c_1L[f_1] + c_2L[f_2] + \dots + c_nL[f_n].$$

5. Show that the following distributive law holds for *linear* operators: if  $L_1, L_2, R_1$ , and  $R_2$  are linear operators, then

$$(L_1 + L_2)(R_1 + R_2) = L_1R_1 + L_1R_2 + L_2R_1 + L_2R_2.$$

Note: Because linear operators don't commute with each other in general (see the next exercise), the formula above is valid *only* with the  $L$ 's in front of the  $R$ 's on the right-hand side of the equation.

6. In class it was stated that general linear differential operators  $L_1, L_2$  do not commute with each other:  $L_1L_2 \neq L_2L_1$ . Part (a) of this problem verifies this statement by giving examples of operators that do not commute. It is still true that *some* linear differential operators commute with each other; you will see examples of this in parts (b) and (c).

(a) Let  $L_1$  be the operator "multiplication by  $p$ ", where  $p$  is a *non-constant* function, and let  $L_2 = D$  (the first-derivative operator). Show that  $L_1L_2 \neq L_2L_1$ .

(b) Let  $L_1$  and  $L_2$  be as in part (a), but this time assume that  $p$  is a *constant* function. Show that in this case  $L_1L_2 = L_2L_1$ .

(c) Let  $a$  and  $b$  be constants, and let  $L_1$  and  $L_2$  be the linear differential operators  $D + a$  and  $D + b$ , respectively. Using part (b) plus Exercise 5 above, show that  $L_1L_2 = L_2L_1$ .

What you have shown in parts (a), (b) and (c) are special cases of a more general principle: all *constant-coefficient* linear differential operators  $a_nD^n + a_{n-1}D^{n-1} + \dots + a_1D + a_0$  commute with each other (here the  $a$ 's are constants), but *non-constant-coefficient* linear differential operators  $p_nD^n + p_{n-1}D^{n-1} + \dots + p_1D + p_0$  (here the  $p$ 's are functions at least one of which

is non-constant) in general do not commute with constant-coefficient operators or with other non-constant-coefficient operators.

7. The method of problems 4.3/38–39 in the book can be adapted to give solutions to several cases of Cauchy-Euler (pronounced “Co-she Oiler”) equations not covered in the book’s exercises. In this problem we consider these other cases.

(a) Fix numbers  $a, b, c$  (with  $a \neq 0$ ) and consider the second-order homogeneous Cauchy-Euler equation

$$ax^2 \frac{d^2y}{dx^2} + bx \frac{dy}{dx} + cy = 0. \quad (1)$$

If we consider this equation on the interval  $\{x < 0\}$ , the substitution  $x = e^t$  cannot be used (why not?). However, using the Chain Rule and the substitution  $x = -u$ , show that (1) for  $x < 0$  is equivalent to the equation

$$au^2 \frac{d^2z}{du^2} + bx \frac{dz}{dx} + cu = 0 \quad (2)$$

for  $u > 0$ , where  $z(u) = y(x) = z(-x)$ . Except for the names of the variables, equations (1) and (2) are the same. Use this to show that if  $\phi(x)$  is a solution of  $ax^2y'' + bxy' + cy = 0$  on the interval  $\{x > 0\}$ , then  $\phi(-x)$  is a solution of the same DE on the interval  $\{x < 0\}$ , and vice-versa. Thus show that if  $y_{\text{gen}}(x)$  is the general solution of (1) on the interval  $\{x > 0\}$ , then  $y_{\text{gen}}(|x|)$  is the general solution of (1) on the interval  $\{x < 0\}$ .

(b) Using the methods of the book’s 4.3/38–39, find the general solution  $y(x)$  of

$$6x^2y'' + xy' + y = 0 \quad (3)$$

on the interval  $\{x > 0\}$ . Then, using part (a) above, find the general solution of (3) on the interval  $\{x < 0\}$ .

(c) Using the methods of the book’s 4.3/38–39, find the general solution  $y(x)$  of

$$x^2y'' + 5xy' + 4y = 0 \quad (4)$$

on the interval  $\{x > 0\}$ . (Remember that your answer must be expressed purely in terms of  $x$ , not partly in terms of  $x$  and partly in terms of  $t$ .) Then, using part (a) above, find the general solution of (4) on the interval  $\{x < 0\}$ .