MAA 4211—OCTOBER 19, 2001

INTERIORS, CLOSURES, AND BOUNDARIES

The facts stated in problems 15-17 on p. 62 of the textbook are of fundamental importance. Here is a summary of these and a few other facts; the proofs are exercises (some of which can be done in one line, and most of which can be done in three lines or less, if you do them in the order listed).

Below, (E, d) always represents an arbitrary metric space, $p \in E$ an arbitrary point, $S \subset E$ an arbitrary subset, and $\mathcal{C}S$ or $\mathcal{C}(S)$ the complement of S in E.

Definitions. Let $S \subset E$. A point p is an *interior point* of S if there exists an open ball centered at p entirely contained in S. The *interior of* S, written Int(S), is defined to be the set of interior points of S. The *closure of* S, written \overline{S} , is defined to be the intersection of all closed sets that contain S. The *boundary of* S, written ∂S , is defined by $\partial S = \overline{S} \cap \overline{CS}$. (In other words, the boundary of a set is the intersection of the closure of the set and the closure of its complement.)

All the facts below are implicitly prefaced with "for all $S \subset E$ ".

Facts about interiors.

(1) $Int(S) \subset S$.

(2) Int(S) is open.

(3) If $U \subset S$ is open, then $U \subset Int(S)$. (Because of this property and properties (1) and (2), Int(S) is often referred to as "the largest open set contained in S".)

(4) S is open iff S = Int(S).

(5) Int(S) is the union of all open sets that are contained in S.

Facts about closures.

(6) \overline{S} is the intersection of all closed sets that contain S (by definition).

- (7) \overline{S} is closed.
- (8) $\overline{S} \supset S$.

(9) If $U \supset S$ is closed, then $U \supset \overline{S}$. (Because of this property and properties (7) and (8), \overline{S} is often referred to as "the smallest closed set containing S".)

(10) S is closed iff $S = \overline{S}$.

(11) $p \in \overline{S}$ iff every open ball centered at p contains a point of S.

(12) $\overline{S} = \mathcal{C}(Int(\mathcal{C}S))$; equivalently, $\mathcal{C}\overline{S} = Int(\mathcal{C}S)$. (In words: the complement of the closure is the interior of the complement.)

(13) \overline{S} is the set of all limits of *E*-convergent sequences of points in *S*. (Here "*E*-convergent" means "convergent in (E, d)".)

Facts about boundaries.

(14) $\partial(\mathcal{C}S) = \partial S.$ (15) $\partial S \subset \overline{S}.$ (16) $E = Int(S) \amalg \partial S \amalg Int(\mathcal{C}S)$. (The disjoint-union symbol \amalg means the almost the same thing as "union", but is applied only to disjoint sets; the statement " $X = Y \amalg Z$ " means that Y and Z are disjoint sets whose union is X.) Remark: everything below is an almost immediate consequence of this fact.

(17) $Int(S) = S - \partial S = \overline{S} - \partial S$. (Recall that A - B denotes the set of points lying in A but not B; equivalently, $A - B = A \cap CB$.)

(18) $\overline{S} = Int(S) \amalg \partial S.$

(19) $\overline{S} = S \cup \partial S$.

(20) S is closed iff $S \supset \partial S$.

(21) S is open iff S and ∂S are disjoint.