

**MAA 4211, Fall 2017—Assignment 5’s non-book problems**

B1. (Strengthening of Rosenlicht problem III.10.) Let  $(p_n)_{n=1}^\infty$  be a convergent sequence in a metric space  $(E, d)$ , let  $R$  be the range of this sequence, and let  $p = \lim_{n \rightarrow \infty} p_n$ . Show that  $R \cup \{p\}$  is a compact subset of  $E$ .

B2. Let  $(E, d)$  be a metric space and let  $S \subset E$ .

(a) Prove that  $\bar{S} = S \cup \{\text{all cluster points of } S\}$ .

(b) Prove that  $\bar{S} = S \cup \{\text{all cluster points of } S \text{ that lie in } \partial S\}$ .

B3. Let  $d_1, d_2$  be equivalent metrics on a set  $E$ . Without using any relations between compactness and sequential compactness, prove that  $(E, d_1)$  is sequentially compact if and only if  $(E, d_2)$  is sequentially compact.

B4. As in previous homework, let  $\mathbf{R}^\infty$  denote the vector space whose elements are real-valued sequences, and let  $\mathbf{R}_b^\infty \subset \mathbf{R}^\infty$  denote the space of bounded real-valued sequences. We will write many elements of  $\mathbf{R}^\infty$  using the notation  $\vec{a}$  to stand for  $(a_i)_{i=1}^\infty$ , the notation  $\vec{b}$  to stand for  $(b_i)_{i=1}^\infty$ , etc. (Thus if we are given an element  $\vec{a} \in \mathbf{R}^\infty$ , it is understood that the notation “ $a_i$ ” means the  $i^{\text{th}}$  term of  $\vec{a}$ .) We often think of elements of  $\mathbf{R}^\infty$  as “infinitely long row-vectors” (i.e. vectors with infinitely many components), and think of the  $i^{\text{th}}$  term of an element of  $\mathbf{R}^\infty$  as the  $i^{\text{th}}$  component of this vector.

The normed vector space  $(\mathbf{R}_b^\infty, \|\cdot\|_\infty)$  is conventionally called  $\ell^\infty(\mathbf{R})$ . As with any normed vector space, when we speak of metric-space properties of  $\ell^\infty(\mathbf{R})$ , the metric is assumed to be the one associated with the given norm (unless otherwise specified); thus the  $\ell^\infty$  metric on  $\mathbf{R}_b^\infty$  is the function  $d : \mathbf{R}_b^\infty \times \mathbf{R}_b^\infty \rightarrow \mathbf{R}$  given by  $d(\vec{a}, \vec{b}) = d_\infty(\vec{a}, \vec{b}) = \sup\{|a_i - b_i| : i \in \mathbf{N}\}$  (exactly the metric in Rosenlicht p. 61/#1b).

Since a sequence in  $\ell^\infty(\mathbf{R})$  is a sequence of sequences, to avoid confusion in this problem we will use a superscript rather than a subscript to label the terms of a sequence in  $\ell^\infty(\mathbf{R})$ ; we will write such a sequence as  $(\vec{a}^{(n)})_{n=1}^\infty$ . Thus the  $n^{\text{th}}$  term in such a sequence is a real-valued sequence  $\vec{a}^{(n)} = (a_i^{(n)})_{i=1}^\infty = (a_1^{(n)}, a_2^{(n)}, a_3^{(n)}, \dots)$ . You may find it helpful to picture such a sequence as an array with infinitely many rows and columns, in which the first row is the sequence  $\vec{a}^{(1)}$ , the second row is the sequence  $\vec{a}^{(2)}$ , etc.:

$$\begin{array}{ccccccc}
 a_1^{(1)} & a_2^{(1)} & a_3^{(1)} & a_4^{(1)} & \dots & & \\
 a_1^{(2)} & a_2^{(2)} & a_3^{(2)} & a_4^{(2)} & \dots & & \\
 a_1^{(3)} & a_2^{(3)} & a_3^{(3)} & a_4^{(3)} & \dots & & \\
 \vdots & \vdots & \vdots & \vdots & \vdots & & 
 \end{array}$$

(a) Let  $(\vec{a}^{(n)})_{n=1}^\infty$  be a Cauchy sequence in  $\ell^\infty(\mathbf{R})$ . Show that for all  $i \in \mathbf{N}$ , the real-valued sequence  $(a_i^{(n)})_{n=1}^\infty$  (the sequence of “ $i^{\text{th}}$  components”—more precisely,  $i^{\text{th}}$  terms—of the  $\vec{a}^{(n)}$ , corresponding to the  $i^{\text{th}}$  column in the diagram above) is a Cauchy

sequence in  $\mathbf{R}$ . Note that in  $(a_i^{(n)})_{n=1}^\infty$ , the index  $i$  is fixed; it is  $n$  that varies:  $(a_i^{(n)})_{n=1}^\infty = (a_i^{(1)}, a_i^{(2)}, a_i^{(3)}, \dots)$ .

(b) Let  $(\vec{a}^{(n)})_{n=1}^\infty$  be as in part (a). Since  $\mathbf{R}$  is complete, for all  $i \in \mathbf{N}$  there exists  $c_i \in \mathbf{R}$  such that  $\lim_{n \rightarrow \infty} a_i^{(n)} = c_i$ . Let  $\vec{c}$  be the sequence  $(c_i)_{i=1}^\infty \in \mathbf{R}^\infty$ —no subscript “ $b$ ”, yet. Show that the sequence  $\vec{c}$  is, in fact, bounded. (So  $\vec{c} \in \mathbf{R}_b^\infty$  after all.)

(c) Let  $(\vec{a}^{(n)})_{n=1}^\infty$  and  $\vec{c}$  be as in part (b). Show that  $(\vec{a}^{(n)})_{n=1}^\infty$  converges in  $\ell^\infty(\mathbf{R})$  to  $\vec{c}$ . (Note: unlike for sequences in  $\mathbf{R}^m$ , this CANNOT be deduced just from the fact that  $(a_i^{(n)})_{n=1}^\infty$  converges to  $c_i$  for all  $i$ ; see part (e) below.) Thus  $\ell^\infty(\mathbf{R})$  is complete.

*Hint:* For  $\epsilon > 0$ , if  $N \in \mathbf{N}$  is as in the Cauchy criterion for the sequence  $(\vec{a}^{(n)})_{n=1}^\infty$  in  $\ell^\infty(\mathbf{R})$ , show that for all  $i \in \mathbf{N}$ , this same  $N$  “works” in the Cauchy criterion for the real-valued sequence  $(a_i^{(n)})_{n=1}^\infty$ . (You probably already did this in part (b).) Then apply a lemma proved in class on 11/8/17 to each sequence  $(a_i^{(n)})_{n=1}^\infty$ .

**Notation for the remaining parts of this problem.** For  $n \in \mathbf{N}$ , let  $\vec{e}^{(n)} \in \mathbf{R}_b^\infty$  be the sequence whose  $n^{\text{th}}$  term is 1 and all of whose other terms are zero (e.g.  $\vec{e}^{(3)} = (0, 0, 1, 0, 0, 0, \dots)$ ).

(d) Show that for all  $i \in \mathbf{N}$ ,  $(e_i^{(n)})_{n=1}^\infty$  converges in  $\mathbf{R}$  to 0.

(e) Let  $\vec{0}$  be the zero element of  $\mathbf{R}_b^\infty$  (the sequence  $(0, 0, 0, 0, \dots)$ ). Compute  $d(\vec{e}^{(n)}, \vec{0})$  for all  $n$ , and use your answer to show that  $(\vec{e}^{(n)})_{n=1}^\infty$  does not converge in  $\ell^\infty(\mathbf{R})$  to  $\vec{0}$ , even though the  $i^{\text{th}}$ -component sequence  $(e_i^{(n)})_{n=1}^\infty$  converges to the  $i^{\text{th}}$  component of  $\vec{0}$  for every  $i$ .

(f) Compute  $d(\vec{e}^{(n)}, \vec{e}^{(m)})$  for all  $m, n \in \mathbf{N}, m \neq n$ . Use your answer to show that no subsequence of  $(\vec{e}^{(n)})_{n=1}^\infty$  can be Cauchy. Use this to deduce that no subsequence of  $(\vec{e}^{(n)})_{n=1}^\infty$  can converge.

Note: since any sequence is trivially a subsequence of itself, the last conclusion implies that  $(\vec{e}^{(n)})_{n=1}^\infty$  does not converge in  $\ell^\infty(\mathbf{R})$  to *anything*, so, in particular, it does not converge to  $\vec{0}$ . But I still want you to do part (e) by the method indicated in part (e).

(g) Use part (f) to deduce that the closed unit ball  $\overline{B}_1(\vec{0}) \subset \ell^\infty(\mathbf{R})$  is not sequentially compact. (Thus this ball is a closed, bounded subset of a complete metric space that generalizes  $(\mathbf{R}^n, d_\infty)$ , but is not compact. Analogs of the Heine-Borel theorem are false in infinite dimensions.)